



The Entiat IMW: Results to Date

Presenter: Pamela Nelle
Collaborators: Shubha Pandit, Tom
Desgroseillier



Entiat Watershed Planning Unit
Entiat, WA
April 6, 2016




Update on Results from Entiat IMW






Experimental Design




Habitat and Fish Monitoring




Habitat Monitoring




Geomorphic Change Detection




Results to Date: Habitat - Geomorphic change



Results to Date: Habitat - Geomorphic change




Results to Date: Habitat




Results to Date: Fish Response

- Results from effectiveness monitoring for spring Chinook and steelhead
 - Abundance
 - Survival
 - Production
 - Productivity
- Annually and by season: winter and summer
- At the IMW (mainstem Entiat and Mad River) scale and valley segment scale


Results to Date: Juvenile Fish Abundance




Results to Date: Annual Survival within IMW




Results to Date: Summer Survival by Valley Segment




Results to Date: Juvenile Fish Winter Survival



Results to Date: Emigration from the Entiat



Results to Date: The Holy Grail = Productivity



Conclusion

- Have a had a real impact on the amount of wood in the river
- Other habitat features important to fish not as great a response or a negative response, but it is early days yet.
- No real change in juvenile Chinook and steelhead parr abundance at the IMW scale
- No real increase in annual survival, but maybe a positive trend for steelhead in the winter
- No increase in survival in the valley segments that were targeted for restoration
- Production trending up for steelhead

Conclusion

- Habitat and fish population restoration not easy to do in the “real world”
- Constraints on what can be done in the instream environment
 - Natural geomorphology
 - Design/permitting
 - Funding
 - Societal
- Continue to strive for as big a restoration impact as possible
- Continue to monitor fish and habitat response to account for many factors involved